

# Pancreatic Cancer: Alarm for Survival

*Felix Aloysius Budimutiar\**, *Marcellus Simadibrata\*\**

\*Department of Internal Medicine, Abdi Waluyo Hospital, Jakarta

\*\* Division of Gastroenterology, Department of Internal Medicine, Faculty of Medicine Universitas Indonesia/Dr. Cipto Mangunkusumo Hospital, Jakarta

## Corresponding author:

*Marcellus Simadibrata. Division of Gastroenterology, Department of Internal Medicine, Dr. Cipto Mangunkusumo General National Hospital. Jl Diponegoro No.71 Jakarta Indonesia. Phone: +62-21-3153957; facsimile: +62-21-3142454. E-mail: prof.marcellus.s@gmail.com*

## ABSTRACT

*Pancreatic cancer is one of the deadliest and a highly aggressive cancer. Its incidence and mortality are highest in developed countries. However, in Asia-Pacific region, the incidence and mortality rate of pancreatic cancer are also on increasing trend nowadays. Pancreatic cancer incidence rates increase with older age, the highest in the 7<sup>th</sup> and 8<sup>th</sup> decades. Clinical presentation of pancreatic cancer are usually non-specific, largely dependent on tumour size and location, with most patients experience symptoms already late in the disease. Computed tomography is considered the method of choice for diagnosis and staging of pancreatic cancer. Management for pancreatic cancer include surgical resection, radiotherapy, chemotherapy. Only surgical resection considered the potentially curative treatment for pancreatic cancer. Unfortunately, a lot of patients present with disease that is not surgically resectable. Prognosis for pancreatic cancer is very poor, despite surgery in resectable patients. The overall five-year survival rate is about less than 5%*

**Keywords:** *pancreatic cancer, diagnosis, management*

## ABSTRAK

*Kanker pankreas merupakan salah satu tipe kanker yang mematikan dan sangat agresif. Insidens dan mortalitas dari kanker pankreas tertinggi adalah dari negara maju. Akan tetapi, di wilayah Asia Pasifik, angka insidens dan mortalitas dari kanker pankreas juga dalam tren meningkat sekarang ini. Insiden kanker pankreas meningkat dengan bertambah tuanya umur, paling tinggi di dekade ke 7 dan 8. Presentasi klinis dari kanker pankreas biasanya non spesifik, tergantung dari ukuran dan lokasi tumor, dengan kebanyakan pasien mengalami gejala sudah di stadium lanjut. Computed tomography dianggap metode pilihan untuk mendiagnosis dan menentukan stadium dari kanker pankreas. Pengobatan kanker pankreas mencakup bedah reseksi, radioterapi dan kemoterapi. Hanya tindakan bedah reseksi yang dianggap berpotensi untuk menyembuhkan kanker pankreas. Sayangnya, kebanyakan pasien datang dengan penyakit kanker pankreas yang sudah tidak bisa direseksi. Prognosis dari kanker pankreas sangat jelek, meskipun pasien dengan kanker pankreas yang bisa direseksi. Angka harapan hidup 5 tahun hanya kurang dari 5%.*

**Kata kunci:** *pancreatic cancer, diagnosis, management*

## INTRODUCTION

According to GLOBOCAN 2012, pancreatic cancer causing more than 331,000 deaths per year, which ranked as seventh leading cause of cancer death in both sexes together. Pancreatic cancer also ranked the 12th most common cancer in the world with 338,000 new cases. The highest incidence and mortality rates of pancreatic cancer are found in developed countries. In Asia Pacific region, the mortality rate of pancreatic cancer are on increasing trend nowadays.<sup>2</sup> The incidence rates of pancreatic cancer for both sexes increase with age, the highest in the 7<sup>th</sup> and 8<sup>th</sup> decades of life. There are many condition that considered as risk factors for pancreatic cancer. The risk factors are divided into modifiable and non-modifiable. Modifiable risk factors include smoking, obesity, alcohol and dietary factors, while non-modifiable risk factors include increasing age, familial cancer syndromes, race, hereditary, diabetes mellitus, and chronic pancreatitis.<sup>3</sup>

The survival rate of pancreatic cancer is very poor; the overall 5-year survival is about less than 5%. Surgical resection is the only potentially curative treatment of choice for pancreatic cancer. The majority of patients are diagnosed at an advanced stage where surgical resection is not an option for treatment, hence the low survival rate. Despite improvements in imaging, surgical techniques and chemotherapy, overall survival has not improved in the past few decades.<sup>1</sup>

## Clinical Presentation

The clinical presentation of patients with pancreatic cancer are usually non-specific, it largely depends on the size and location of the tumor. Pancreatic cancer location approximately 60-70% in the head of pancreas, 20-25% in the body and tail of pancreas, dan the remaining 10-20% diffusely involve the pancreas. Symptoms of pancreatic cancer include jaundice, abdominal pain, weight loss, steatorrhoea, and new onset diabetes.<sup>4</sup>

## Diagnostics

Pancreas protocol CT is the gold standard for diagnosis and staging pancreatic cancer. The pancreatic protocol CT consists of dual-phase scanning using intravenous and oral contrast agents. It is the most widely used modality for diagnosis and staging of pancreatic cancer due to the fairly wide availability. CT can be used to assess potential vascular involvement and metastases.<sup>5,6</sup> Although the widely used of CT for diagnosis of pancreatic cancer, MRI can also play an

important role. MRI have more superior soft tissue contrast that can detect small, non-countour deforming tumors which may not be seen on CT. MRI and CT have similar performance for staging of pancreatic cancer. In some cases, MRI improves detection of hepatic metastases and characterization of CT-indeterminate liver lesions.<sup>7</sup>

Another modality used in the diagnosis of pancreatic cancer is endoscopic ultrasound (EUS). EUS with Fine Needle Aspiration (FNA) can help secure tissue for cytologic diagnosis which made it essential for diagnosing pancreatic cancer. EUS can provide staging of suspected or proven pancreatic cancer, and cytological/histological proof of unresectable pancreatic cancer.<sup>8</sup> Although EUS is a minimally invasive procedure, it is considered a safe procedure with low complication rates. Complications of EUS procedure include bleeding, pancreatitis, perforation, and rarely tumor seeding after EUS-FNA.<sup>9</sup>

## Staging

The American Joint Committee on Cancer (AJCC) 8<sup>th</sup> edition staging system for pancreatic cancer is shown Table 1. This new staging system different from the previous 7<sup>th</sup> edition. The T category is revised from descriptive to size-based definitions. Another change, resectability is removed from the T4 category because the definition of resectability is not consistent among institutions and evolves with advances in surgical technique.<sup>10</sup>

**Table 1. American Joint Committee on Cancer (AJCC) 8<sup>th</sup> edition staging system for pancreatic cancer**

|   |   |       |    |
|---|---|-------|----|
| Tumor (T)   |   |       |    |
| TX  | Primary tumor cannot be assessed  |       |    |
| T0  | No evidence of primary tumor  |       |    |
| Tis   | Carcinoma in situ (includes PanIN III)  |       |    |
| T1  | Maximum tumor diameter ≤ 2 cm   |       |    |
| T2  | Maximum tumor diameter >2, ≤4 cm  |       |    |
| T3  | Maximum tumor diameter >4 cm  |       |    |
| T4  | Tumor involves the celiac axis, common hepatic artery or the superior mesenteric artery |       |    |
| Lymph node metastases (N)                         |   |       |    |
| N0  | No regional lymph node metastasis   |       |    |
| N1  | Metastasis in 1–3 regional lymph nodes  |       |    |
| N2  | Metastasis in ≥4 regional lymph nodes   |       |    |
| Distant metastases (M)                            |   |       |    |
| M0  | No distant metastasis   |       |    |
| M1  | Distant metastasis  |       |    |
| American Joint Committee on Cancer (AJCC) staging |   |       |    |
| Stage IA  | T1  | N0    | M0 |
| Stage IB  | T2  | N0    | M0 |
| Stage IIA   | T3  | N0    | M0 |
| Stage IIB   | T1-3  | N1    | M0 |
| Stage III   | Any T   | N2    | M0 |
|   | T4  | Any N | M0 |
| Stage IV  | Any T   | Any N | M1 |

Reprinted from American Joint Committee on Cancer (AJCC) 8<sup>th</sup> edition

## TREATMENT

### Resectable

Surgical resection is a potentially curative treatment for pancreatic cancer, but only 15–20% of patients present with a resectable disease.<sup>11</sup> Patients who have localized disease with no metastases, good performance status and no significant comorbidities are a good candidate for surgical resection. The classic surgery for resection is a pancreaticoduodenectomy, also known as Whipple procedure.<sup>12</sup> Pancreatic cancer patients who undergo surgical resection is recommended to be given adjuvant chemotherapy.<sup>13</sup> ESPAC-4 trial showed that patients with resected pancreatic cancer who were given adjuvant combination of gemcitabine and capecitabine had improve overall survival compared with only gemcitabine (OS: 28 months vs. 25.5 months; HR for death 0.82; 95% CI 0.68–0.98).<sup>14</sup> There is still no recommendation to giving neoadjuvant chemotherapy in resectable pancreatic cancer because no significant improvement in overall survival.<sup>15</sup>

### Borderline Resectable

Borderline resectable pancreatic cancer have a higher risk for positive margin after resection. Therefore, neoadjuvant therapy recommended to be given before surgical resection of borderline resectable pancreatic cancer. A study using FOLFIRINOX followed by capecitabine based chemoradiation for borderline resectable pancreatic cancer showed a successful pancreatectomy (68%) with negative margins resection rates (93%).<sup>16</sup>

### Locally Advanced

Patients with locally advanced pancreatic cancer who underwent surgical resection usually had a poor prognosis due to high rates of margin positivity after surgical resection. Systemic chemotherapy is the preferred treatment for locally advanced pancreatic cancer. A systematic review and meta analysis study using FOLFIRINOX as systematic chemotherapy showed a median overall survival longer (24.2 months) compared to gemcitabine (6-13 months). The study also showed about 28% patients were able to underwent surgical resection of tumour and 78% of them had negative margin resection.<sup>17</sup> Chemotherapy in this situation is never curative, and its palliative benefit must be balanced against its potential toxic side effects. Despite treatment, the survival outcomes remain poor.<sup>18</sup>

## Endoscopic Ultrasound (EUS) Interventional Therapy

EUS interventional therapy in pancreatic cancer management is emerging. Many applications using EUS include fiducials placement, brachytherapy, fine needle injection (FNI) and radiofrequency ablation (RFA). Fiducial are inert radiopaque spheres, coils, or cylindricals that are implanted inside or adjacent to the tumor in order to aid image-guided radiation therapy (IGRT). EUS-guided fiducials placement first reported by Pishvaian *et al* in case series of 13 patients where technical success achieved in 94% patients.<sup>19</sup> Followed by several case series with technical success more than 90%.<sup>20-22</sup>

EUS-brachytherapy apply the insertion of a radioactive seed (iodine-125, iridium-192, and palladium-103) directly into the pancreatic tumor. Brachytherapy still need more larger studies for evaluation. Studies done showed EUS-brachytherapy safe, feasible and have benefit in pain control.<sup>23-25</sup>

EUS-FNI apply direct injection of therapeutic agents into pancreatic tumors. EUS-FNI is safe and feasible, but not yet show long term efficiency.<sup>26-27</sup> EUS-RFA role in management of pancreatic cancer had been showed in recent studies. A study by Song *et al* demonstrated safety and feasibility of EUS-RFA in patients with unresectable pancreatic cancer. Technical success in this study was achieved 100% however overall survival can't be assessed.<sup>28</sup>

## CONCLUSION

Pancreatic cancer still considered one of the deadliest cancer types. Survival rate of this disease is very poor. There are many diagnostics modality for diagnosing pancreatic cancer but early diagnosis of pancreatic cancer is still difficult. One of the factor is because patients only experience symptoms of pancreatic cancer late in the course of the disease. Despite treatment, pancreatic cancer high aggressiveness and resistance to modern chemo-radiotherapy are associated with low rates of long-term survival. Studies using EUS as interventional therapy showed promising results. More larger studies are needed to establish the role of these procedures in the treatment of pancreatic cancer. There are no screening recommendations for pancreatic cancer, so primary prevention is of utmost importance.

## REFERENCES

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-86.
2. Pourhoseingholi MA, Ashtari S, Hajizadeh N, Fazeli Z, Zali MR. Systematic review of pancreatic cancer epidemiology in Asia-Pacific Region: major patterns in GLOBOCAN 2012. *Gastroenterol Hepatol Bed Bench* 2017;10:245-57.
3. Midha S, Chawla S, Garg PK. Modifiable and non-modifiable risk factors for pancreatic cancer: A review. *Cancer Lett* 2016;381:269-77.
4. Ducreux M, Cuhna AS, Caramella C, Hollebecque A, Burtin P. Cancer of the pancreas: ESMO clinical practice, guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2015; 26:56-68.
5. Al-Hawary M. Role of imaging in diagnosing and staging pancreatic cancer. *J Natl Compr Canc Netw* 2016;14:678-80.
6. Tempero MA, Malafa MP, Al-Hawary M, Asbun H, Bain A, Behrman SW, et al. Pancreatic Adenocarcinoma, Version 2, 2017. *J Natl Compr Canc Netw* 2017;15:1028-61.
7. Jha P, Yeh BM, Zagoria R, Collison E, Wang ZJ. The role of MR imaging in pancreatic cancer. *Magn Reson Imaging Clin N Am* 2018;26:363-73.
8. Goncalves B, Soares JB, Bastos P. Endoscopic ultrasound in the diagnosis and staging of pancreatic cancer. *Port J Gastroenterol* 2015;22:161-71.
9. Gonzalo-Marin J, Vila JJ, Pirez-Miranda M. Role of endoscopic ultrasound in the diagnosis of pancreatic cancer. *World J Gastrointest Oncol.* 2014;6:360-8.
10. Chun YS, Pawlik TM, Vauthey JN. 8<sup>th</sup> edition of the AJCC cancer staging manual: pancreas and hepatobiliary cancers. *Ann Surg Oncol* 2017;25:845-7.
11. Hackert T, Ulrich A, Buchler MW. Can neoadjuvant therapy in pancreatic cancer increase the pool of patients eligible for pancreaticoduodenectomy? *Adv Surg* 2017;51:1-10.
12. Saraee A, Vahedian-Arkadani J, Saraee E, Pakzad R, Wadji MB. Whipple procedure: a review of a 7 year clinical experience in a referral center for hepatobiliary and pancreas diseases. *World J Surg Oncol.* 2015;13: 98.
13. Casadei R, Di Marco M, Ricci C, Santini D, Serra C, Calculli L, et al. Neoadjuvant chemoradiotherapy and surgery versus surgery alone in resectable pancreatic cancer: a single-center prospective, randomized, controlled trial which failed to achieve accrual targets. *J Gastrointest Surg* 2015;19:1802-12.
14. Neoptolemos JP, Palmer DH, Ghaneh P, Psarelli EE, Valle JW, Halloran CM, et al. Comparison of adjuvant gemcitabine and capecitabine with gemcitabine monotherapy in patients with resected pancreatic cancer (ESPAC-4): A multicentre, open-label, randomised, phase 3 trial. *Lancet.* 2017;389:1011-24.
15. Rahman SH, Urquhart R, Molinari M. Neoadjuvant therapy for resectable pancreatic cancer. *World J Gastrointest Oncol.* 2017;9:457-65.
16. Katz MH, Shi Q, Ahmad SA, Herman JM, Marsh RW, Collisson E, et al. Preoperative modified folfox treatment followed by capecitabine-based chemoradiation for borderline resectable pancreatic cancer: Alliance for clinical trials in oncology trial a021101. *JAMA Surg.* 2016;151:e161137.
17. Suker M, Beumer BR, Sadot E, Marthey L, Faris JE, Mellon EA, et al. Folfirinox for locally advanced pancreatic cancer: A systematic review and patient-level meta-analysis. *Lancet Oncol.* 2016;17:801-10.
18. Sohal DPS, Kennedy EB, Khorana A, Copur MS, Crane CH, Laguna IG, et al. Metastatic pancreatic cancer: ASCO Clinical Practice Guideline Update. *J Clin Oncol* 2018;36:2545-56.
19. Pishvaian AC, Collins B, Gagnon G, Ahlawat S, Haddad NG. EUS-guided fiducial placement for Cyber Knife radiotherapy of mediastinal and abdominal malignancies. *Gastrointest Endosc* 2006;64:412-17.
20. Sanders MK, Moser AJ, Khalid A, Fasanella KE, Zeh HJ, Burton S, et al. EUS-guided fiducial placement for stereotactic body radiotherapy in locally advanced and recurrent pancreatic cancer. *Gastrointest Endosc* 2010;71:1178-84.
21. Park WG, Yan BM, Schellenberg D, Kim J, Chang DT, Koong A, et al. EUS-guided gold fiducial insertion for image-guided radiation therapy of pancreatic cancer: 50 successful cases without fluoroscopy. *Gastrointest Endosc* 2010;71:513-18.
22. Khashab MA, Kim KJ, Tryggstad EJ, Wild AT, Roland T, Singh VK, et al. Comparative analysis of traditional and coiled fiducials implanted during EUS for pancreatic cancer patients receiving stereotactic body radiation therapy. *Gastrointest Endosc* 2012;76:962-71.
23. Sun S, Xu H, Xin J, Liu J, Guo Q, Li S. Endoscopic ultrasound-guided interstitial brachytherapy of unresectable pancreatic cancer: results of a pilot trial. *Endoscopy* 2006;38:399-403.
24. Jin Z, Du Y, Li Z, Jiang Y, Chen J, Liu Y. Endoscopic ultrasonography-guided interstitial implantation of iodine 125-seeds combined with chemotherapy in the treatment of unresectable pancreatic carcinoma: a prospective pilot study. *Endoscopy* 2008;40:314-20.
25. Du Y, Jin Z, Meng H, Zou D, Chen J, Liu Y, et al. Long-term effect of gemcitabine-combined endoscopic ultrasonography-guided brachytherapy in pancreatic cancer. *J Interv Gastroenterol* 2013;3:8-24.
26. Chang KJ, Nguyen PT, Thompson JA, Kurosaki TT, Casey LR, Leung EC, et al. Phase I clinical trial of allogeneic mixed lymphocyte culture (cytoimplant) delivered by endoscopic ultrasound-guided fine-needle injection in patients with advanced pancreatic carcinoma. *Cancer* 2000;88:1325-35.
27. Hecht JR, Bedford R, Abbruzzese JL, Lahoti S, Reid TR, Soetikno RM, et al. A phase I/II trial of intratumoral endoscopic ultrasound injection of ONYX-015 with intravenous gemcitabine in unresectable pancreatic carcinoma. *Clin Cancer Res* 2003;9:555-561.
28. Song TJ, Seo DW, Lakhtakia S, Reddy N, Oh DW, Park do H, et al. Initial experience of EUS-guided radiofrequency ablation of unresectable pancreatic cancer. *Gastrointest Endosc* 2016;83:440-3.